

**Correlation of
AP[®] Chemistry Standards to
Chemistry: A Molecular Approach, 6e AP Edition
(Tro)**



**Correlation of AP[®] Chemistry Standards to *Chemistry: A Molecular Approach*
6th Edition, AP Edition, (Tro)**

Unit 1: Atomic Structure and Properties

AP Topic	Learning Objective	Essential Knowledge	Textbook Chapter & Section and Page Reference
1.1 Moles and Molar Mass	1.1.A Calculate quantities of a substance or its relative number of particles using dimensional analysis and the mole concept	1.1.A.1 One cannot count particles directly while performing laboratory work. Thus, there must be a connection between the masses of substances reacting and the actual number of particles undergoing chemical changes.	Ch. 3, Sec. 3.1, 3.3–3.4; Pgs 91-92, 94-99
		1.1.A.2 Avogadro's number ($N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$) provides the connection between the number of moles in a pure sample of a substance and the number of constituent particles (or formula units) of that substance	Ch. 6, Sec 6.3; Pgs 217, 221-223
		1.1.A.3 Expressing the mass of an individual atom or molecule in atomic mass units (amu) is useful because the average mass in amu of one particle (atom or molecule) or formula unit of a substance will always be numerically equal to the molar mass of that substance in grams. Thus, there is a quantitative connection between the mass of a substance and the number of particles that the substance contains. EQN: $n = m/M$	Ch. 2, Sec 2.8; Pgs 69-72



AP Topic	Learning Objective	Essential Knowledge	Textbook Chapter & Section and Page Reference
1.2 Mass Spectra of Elements	1.2.A Explain the quantitative relationship between the mass spectrum of an element and the masses of the element's isotopes.	1.2.A.1 The mass spectrum of a sample containing a single element can be used to determine the identity of the isotopes of that element and the relative abundance of each isotope in nature	Ch. 2, Sec. 2.8; Pgs 69-72
		1.2.A.2 The average atomic mass of an element can be estimated from the weighted average of the isotopic masses using the mass of each isotope and its relative abundance	Ch. 2, Sec. 2.8; Pgs 69-72
1.3 Elemental Composition of Pure Substances	1.3.A Explain the quantitative relationship between the elemental composition by mass and the empirical formula of a pure substance	1.3.A.1 Some pure substances are composed of individual molecules, while others consist of atoms or ions held together in fixed proportions as described by a formula unit.	Ch. 3, Sec. 3.9-3.10; Pgs 113-122
		1.3.A.2 According to the law of definite proportions, the ratio of the masses of the constituent elements in any pure sample of that compound is always the same.	Ch. 2, Sec. 2.3; Pgs 52-53
		1.3.A.3 The chemical formula that lists the lowest whole number ratio of atoms of the elements in a compound is the empirical formula.	Ch. 3, Sec. 3.3, 94-96, Sec. 3.10; Pgs 118-121, Sec 3.9, Pgs 116



1.4 Composition of Mixtures	1.4.A Explain the quantitative relationship between the elemental composition by mass and the composition of substances in a mixture.	1.4.A.1 Pure substances contain atoms, molecules, or formula units of a single type. Mixtures contain atoms, molecules, or formula units of two or more types, whose relative proportions can vary.	Ch. 1, Sec. 1.3-1.4; Pgs 5-11
		1.4.A.2 Elemental analysis can be used to determine the relative numbers of atoms in a substance and to determine its purity.	Ch. 1, Sec. 1.3-1.4; Pgs 5-11
1.5 Atomic Structure and Electron Configuration	1.5.A Represent the ground-state electron configuration of an atom of an element or its ions using the Aufbau principle.	1.5.A.1 The atom is composed of negatively charged electrons and a positively charged nucleus that is made of protons and neutrons.	Ch. 9, Sec. 9.3-9.4; Pgs 361-372
		1.5.A.2 Coulomb's law is used to calculate the force between two charged particles. EQN: $F_{coulombic} \propto \frac{q_1 q_2}{r^2}$	Ch. 9, Sec. 9.3-9.4; Pgs 363-366
		1.5.A.3 In atoms and ions, the electrons can be thought of as being in "shells (energy levels)" and "subshells (sublevels)," as described by the ground-state electron configuration. Inner electrons are called core electrons, and outer electrons are called valence electrons. The electron configuration is explained by quantum mechanics, as delineated in the Aufbau principle and exemplified in the periodic table of the elements.	Ch. 9, Sec. 9.3-9.4; Pgs 363-366



		1.5.A.4 The relative energy required to remove an electron from different subshells of an atom or ion or from the same subshell in different atoms or ions (ionization energy) can be estimated through a qualitative application of Coulomb's law. This energy is related to the distance from the nucleus and the effective (shield) charge of the nucleus.	Ch. 9, Sec. 9.7; Pgs 379-386. Ch. 9, Sec. 9.3-9.4; Pgs 363-366
1.6 Photoelectron Spectroscopy (PES)	1.6.A Explain the relationship between the photoelectron spectrum of an atom or ion and: i. The ground-state electron configuration of the species. ii. The interactions between the electrons and the nucleus.	1.6.A.1 The energies of the electrons in a given shell can be measured experimentally with photoelectron spectroscopy (PES). The position of each peak in the PES spectrum is related to the energy required to remove an electron from the corresponding subshell, and the relative height of each peak is (ideally) proportional to the number of electrons in that subshell.	Not explicitly covered in the text
1.7 Periodic Trends	1.7.A Explain the relationship between trends in atomic properties of elements and electronic structure and periodicity.	1.7.A.1 The organization of the periodic table is based on patterns of recurring properties of the elements, which are explained by patterns of ground-state electron configurations and the presence of completely or partially filled shells (and subshells) of electrons in atoms.	Ch. 9, Sec. 9.2, 9.5-9.9; 360-361, 373-391
		1.7.A.2 Trends in atomic properties within the periodic table (periodicity) can be predicted by the position of the	Ch. 9, Sec. 9.3-9.4; Pgs 363-366. Ch. 10, Sec 10.4; Pgs 405-



		element on the periodic table and qualitatively understood using Coulomb's law, the shell model, and the concepts of shielding and effective nuclear charge. These properties include: i. Ionization energy ii. Atomic and ionic radii iii. Electron affinity iv. Electronegativity.	411, Ch 13, Sec 13.5; Pgs 566-567. Ch. 9, Sec. 9.7; Pgs 379-386.
		1.7.A.3 The periodicity (in 1.7.A.2) is useful to predict/ estimate values of properties in the absence of data.	Ch. 9, Sec. 9.3-9.4; Pgs 363-366. Ch. 10, Sec 10.4; Pgs 405-411, Ch 13, Sec 13.5; Pgs 566-567. Ch. 9, Sec. 9.7; Pgs 379-386.
1.8 Valence Electrons and Ionic Compounds	1.8.A Explain the relationship between trends in the reactivity of elements and periodicity.	1.8.A.1 The likelihood that two elements will form a chemical bond is determined by the interactions between the valence electrons and nuclei of elements.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411
		1.8.A.2 Elements in the same column of the periodic table tend to form analogous compounds.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411
		1.8.A.3 Typical charges of atoms in ionic compounds are governed by the number of valence electrons and predicted by their location on the periodic table.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411



Unit 2: Compound Structure and Properties

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
2.1 Types of Chemical Bonds	2.1.A Explain the relationship between the type of bonding and the properties of the elements participating in the bond.	2.1.A.1 Electronegativity values for the representative elements increase going from left to right across a period and decrease going down a group. These trends can be understood qualitatively through the electronic structure of the atoms, the shell model, and Coulomb's law.	Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
		2.1.A.2 Valence electrons shared between atoms of similar electronegativity constitute a nonpolar covalent bond. For example, bonds between carbon and hydrogen are effectively nonpolar even though carbon is slightly more electronegative than hydrogen.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
		2.1.A.3 Valence electrons shared between atoms of unequal electronegativity constitute a polar covalent bond. i. The atom with a higher electronegativity will develop a partial negative charge relative to the other atom in the bond. ii. In single bonds, greater differences in electronegativity lead to greater bond dipoles.	Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
		iii. All polar bonds have some ionic character, and the difference between ionic and covalent bonding is not distinct but rather a continuum.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		2.1.A.4 The difference in electronegativity is not the only factor in determining if a bond should be designated as ionic or covalent. Generally, bonds between a metal and nonmetal are ionic, and bonds between two nonmetals are covalent. Examination of the properties of a compound is the best way to characterize the type of bonding.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
		2.1.A.5 In a metallic solid, the valence electrons from the metal atoms are considered to be delocalized and not associated with any individual atom.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
2.2 Intramolecular Force and Potential Energy	2.2.A Represent the relationship between potential energy and distance between atoms, based on factors that influence the interaction strength.	2.2.A.1 A graph of potential energy versus the distance between atoms (internuclear distance) is a useful representation for describing the interactions between atoms. Such graphs illustrate both the equilibrium bond length (the separation between atoms at which the potential energy is lowest) and the bond energy (the energy required to separate the atoms).	Ch. 10, Sec. 10.10; Pgs 430-432
		2.2.A.2 In a covalent bond, the bond length is influenced by both the size of the atom's core and the bond order (i.e., single, double, triple). Bonds with a higher order are	Ch. 10, Sec. 10.5; Pgs 412-413



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		shorter and have larger bond energies.	
		2.2.A.3 Coulomb's law can be used to understand the strength of interactions between cations and anions. i. Because the interaction strength is proportional to the charge on each ion, larger charges lead to stronger interactions. ii. Because the interaction strength increases as the distance between the centers of the ions (nuclei) decreases, smaller ions lead to stronger interactions.	Ch. 9, Sec. 9.3-9.4; Pgs 363-366. Ch. 10, Sec 10.4; Pgs 405-411, Ch 13, Sec 13.5; Pgs 566-567
2.3 Structure of Ionic Solids	2.3.A Represent an ionic solid with a particulate model that is consistent with Coulomb's law and the properties of the constituent ions.	2.3.A.1 The cations and anions in an ionic crystal are arranged in a systematic, periodic 3-D array that maximizes the attractive forces among cations and anions while minimizing the repulsive forces.	Ch. 10, Sec 10.4; Pgs 405-411, Ch 13, Sec 13.5; Pgs 566-567
2.4 Structure of Metals and Alloys	2.4.A Represent a metallic solid and/or alloy using a model to show essential characteristics of the structure and interactions present in the substance.	2.4.A.1 Metallic bonding can be represented as an array of positive metal ions surrounded by delocalized valence electrons (i.e., a "sea of electrons").	Ch 10, Sec. 10.2; Pgs 402-403, Ch 13, Sec 13.4; Pgs 562-565, Ch 25, Sec. 25.4; 1138-1142
		2.4.A.2 Interstitial alloys form between atoms of significantly different radii, where the smaller	Ch. 25, Sec 25.4; pg: 1141



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		atoms fill the interstitial spaces between the larger atoms (e.g., with steel in which carbon occupies the interstices in iron).	
		2.4.A.3 Substitutional alloys form between atoms of comparable radius, where one atom substitutes for the other in the lattice. (e.g., in certain brass alloys, other elements, usually zinc, substitute for copper.)	Ch. 25, Sec 25.4; pg: 1141
2.5 Lewis Diagrams	2.5.A Represent a molecule with a Lewis diagram.	2.5.A.1 Lewis diagrams can be constructed according to an established set of principles.	Ch. 10, Sec. 10.1–10.3, 10.7; Pgs 401-404, 418-419
2.6 Resonance and Formal Charge	2.6.A Represent a molecule with a Lewis diagram that accounts for resonance between equivalent structures or that uses formal charge to select between nonequivalent structures.	2.6.A.1 In cases where more than one equivalent Lewis structure can be constructed, resonance must be included as a refinement to the Lewis structure. In many such cases, this refinement is needed to provide qualitatively accurate predictions of molecular structure and properties.	Ch. 10, Sec. 10.8; Pgs 420-424
		2.6.A.2 The octet rule and formal charge can be used as criteria for determining which of several possible valid Lewis diagrams provides the best model for predicting molecular structure and properties.	Ch. 10, Sec. 10.9; Pgs 425-428



		2.6.A.3 As with any model, there are limitations to the use of the Lewis structure model, particularly in cases with an odd number of valence electrons.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
2.7 VSEPR and Hybridization	2.7.A Based on the relationship between Lewis diagrams, VSEPR theory, bond orders, and bond polarities: i. Explain structural properties of molecules. ii. Explain electron properties of molecules.	2.7.A.1 VSEPR theory uses the Coulombic repulsion between electrons as a basis for predicting the arrangement of electron pairs around a central atom.	Ch. 11, Sec. 11.1-11.4, 11.7; Pgs 445-457, 465-477
		2.7.A.2 Both Lewis diagrams and VSEPR theory must be used for predicting electronic and structural properties of many covalently bonded molecules and polyatomic ions, including the following: i. Molecular geometry (linear, trigonal planar, tetrahedral, trigonal pyramidal, bent, trigonal bipyramidal, seesaw, T-shaped, octahedral, square pyramidal, square planar) ii. Bond angles iii. Relative bond energies based on bond order iv. Relative bond lengths (multiple bonds, effects of atomic radius) v. Presence of a dipole moment vi. Hybridization of valence orbitals for atoms within a molecule or polyatomic ion	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417



		2.7.A.3 The terms “hybridization” and “hybrid atomic orbital” are used to describe the arrangement of electrons around a central atom. When the central atom is sp hybridized, its ideal bond angles are 180° ; for sp^2 hybridized atoms the bond angles are 120° ; and for sp^3 hybridized atoms the bond angles are 109.5° .	Ch. 11, Sec. 11.1-11.4, 11.7; Pgs 445-457, 465-477
		2.7.A.4 Bond formation is associated with overlap between atomic orbitals. In multiple bonds, such overlap leads to the formation of both sigma and pi bonds. The overlap is stronger in sigma than pi bonds, which is reflected in sigma bonds having greater bond energy than pi bonds. The presence of a pi bond also prevents the rotation of the bond and leads to geometric isomers.	Ch. 11, Sec. 11.1-11.4, 11.7; Pgs 445-457, 465-477

Unit 3: Properties of Substances and Mixtures

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
3.1 Intermolecular and Interparticle Forces	3.1.A Explain the relationship between the chemical structures of molecules and the relative strength of their intermolecular forces when: i. The molecules are of the same chemical species. ii. The molecules are	3.1.A.1 London dispersion forces are a result of the Coulombic interactions between temporary, fluctuating dipoles. London dispersion forces are often the strongest net intermolecular force between large molecules. i. Dispersion forces increase with increasing contact area between molecules and with increasing polarizability of the molecules. ii. The polarizability of a molecule	Ch. 12, Sec. 12.2-12.3; Pgs 504-516



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	of two different chemical species	increases with an increasing number of electrons in the molecule and the size of the electron cloud. It is enhanced by the presence of pi bonding. iii. The term "London dispersion forces" should not be used synonymously with the term "van der Waals forces."	
		3.1.A.2 The dipole moment of a polar molecule leads to additional interactions with other chemical species. i. Dipole-induced dipole interactions are present between a polar and nonpolar molecule. These forces are always attractive. The strength of these forces increases with the magnitude of the dipole of the polar molecule and with the polarizability of the nonpolar molecule. ii. Dipole-dipole interactions are present between polar molecules. The interaction strength depends on the magnitudes of the dipoles and their relative orientation. Interactions between polar molecules are typically greater than those between nonpolar molecules of comparable size because these interactions act in addition to London dispersion forces. iii. Ion-dipole forces of attraction are present between ions and polar molecules. These tend to be	Ch. 12, Sec. 12.2-12.3; Pgs 504-516



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		stronger than dipole-dipole forces.	
		3.1.A.3 The relative strength and orientation dependence of dipole-dipole and ion-dipole forces can be understood qualitatively by considering the sign of the partial charges responsible for the molecular dipole moment, and how these partial charges interact with an ion or with an adjacent dipole.	Ch. 12, Sec. 12.2-12.3; Pgs 504-516
		3.1.A.4 Hydrogen bonding is a strong type of intermolecular interaction that exists when hydrogen atoms covalently bonded to the highly electronegative atoms (N, O, and F) are attracted to the negative end of a dipole formed by the electronegative atom (N, O, and F) in a different molecule, or a different part of the same molecule.	Ch. 12, Sec. 12.2-12.3; Pgs 504-516
		3.1.A.5 In large biomolecules, noncovalent interactions may occur between different molecules or between different regions of the same large biomolecule.	Ch. 12, Sec. 12.2-12.3; Pgs 504-516
3.2 Properties of Solids	3.2.A Explain the relationship among the macroscopic properties of a substance, the particulate-level structure of the	3.2.A.1 Many properties of liquids and solids are determined by the strengths and types of intermolecular forces present. Because intermolecular interactions are overcome completely when a substance	Ch. 12, Sec. 12.2, 12.4-12.6; Pgs 504-506, 517-531



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	substance, and the interactions between these particles.	vaporizes, the vapor pressure and boiling point are directly related to the strength of those interactions. Melting points also tend to correlate with interaction strength, but because the interactions are only rearranged, in melting, the relations can be more subtle.	
		3.2.A.2 Particulate-level representations, showing multiple interacting chemical species, are a useful means to communicate or understand how intermolecular interactions help to establish macroscopic properties.	Ch. 12, Sec. 12.2, 12.4–12.6; Pgs 504-506, 517-531
		3.2.A.3 Due to strong interactions between ions, ionic solids tend to have low vapor pressures, high melting points, and high boiling points. They tend to be brittle due to the repulsion of like charges caused when one layer slides across another layer. They conduct electricity only when the ions are mobile, as when the ionic solid is melted (i.e., in a molten state) or dissolved in water or another solvent.	Ch. 12, Sec. 12.2, 12.4–12.6; Pgs 504-506, 517-531
		3.2.A.4 In covalent network solids, the atoms are covalently bonded together into a three dimensional network (e.g., diamond) or layers of two-dimensional networks (e.g., graphite). These are only formed from nonmetals and metalloids: elemental (e.g., diamond, graphite) or binary compounds (e.g., silicon	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		dioxide and silicon carbide). Due to the strong covalent interactions, covalent solids have high melting points. Three-dimensional network solids are also rigid and hard, because the covalent bond angles are fixed. However, graphite is soft because adjacent layers can slide past each other relatively easily.	
		3.2.A.5 Molecular solids are composed of distinct, individual units of covalently-bonded molecules attracted to each other through relatively weak intermolecular forces. Molecular solids generally have a low melting point because of the relatively weak intermolecular forces present between the molecules. They do not conduct electricity because their valence electrons are tightly held within the covalent bonds and the lone pairs of each constituent molecule. Molecular solids are sometimes composed of very large molecules or polymers.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
		3.2.A.6 Metallic solids are good conductors of electricity and heat, due to the presence of free valence electrons. They also tend to be malleable and ductile, due to the ease with which the metal cores can rearrange their structure. In an interstitial alloy, interstitial atoms tend to make the lattice more rigid, decreasing malleability and ductility. Alloys	Ch. 25, Sec 25.4; pg: 1141



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		typically retain a sea of mobile electrons and so remain conducting.	
		3.2.A.7 In large biomolecules or polymers, noncovalent interactions may occur between different molecules or between different regions of the same large biomolecule. The functionality and properties of such molecules depend strongly on the shape of the molecule, which is largely dictated by noncovalent interactions.	Text does not thoroughly explain the standard.
3.3 Solids, Liquids, and Gases	3.3.A Represent the differences between solid, liquid, and gas phases using a particulate level model.	3.3.A.1 Solids can be crystalline, where the particles are arranged in a regular three-dimensional structure, or they can be amorphous, where the particles do not have a regular, orderly arrangement. In both cases, the motion of the individual particles is limited, and the particles do not undergo overall translation with respect to each other. The structure of the solid is influenced by interparticle interactions and the ability of the particles to pack together.	Ch. 12, Sec. 12.7-12.9; Pgs 532-539
		3.3.A.2 The constituent particles in liquids are in close contact with each other, and they are continually moving and colliding. The arrangement and movement of particles are influenced by the	Ch. 12, Sec. 12.7-12.9; Pgs 532-539



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		nature and strength of the forces (e.g., polarity, hydrogen bonding, and temperature) between the particles.	
		3.3.A.3 The solid and liquid phases for a particular substance typically have similar molar volume because, in both phases, the constituent particles are in close contact at all times.	Ch. 12, Sec. 12.7-12.9; Pgs 532-539
		3.3.A.4 In the gas phase, the particles are in constant motion. Their frequencies of collision and the average spacing between them are dependent on temperature, pressure, and volume. Because of this constant motion, and minimal effects of forces between particles, a gas has neither a definite volume nor a definite shape.	Ch. 12, Sec. 12.7-12.9; Pgs 532-539
3.4 Ideal Gas Law	3.4.A Explain the relationship between the macroscopic properties of a sample of gas or mixture of gases using the ideal gas law.	3.4.A.1 The macroscopic properties of ideal gases are related through the ideal gas law: EQN: $PV = nRT$	Ch. 6, Sec. 6.2-6.6; Pgs 214-236
		3.4.A.2 In a sample containing a mixture of ideal gases, the pressure exerted by each component (the partial pressure) is independent of the other components. Therefore, the partial pressure of a gas within the mixture is proportional to its mole	Ch. 6, Sec. 6.2-6.6; Pgs 214-236



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		fraction (X), and the total pressure of the sample is the sum of the partial pressures. EQN: $P_A = P_{total} \times X_A$, where X_A = moles A/total moles; EQN: $P_{total} = P_A + P_B + P_C + \dots$	
		3.4.A.3 Graphical representations of the relationships between P , V , T , and n are useful to describe gas behavior.	Ch. 6, Sec. 6.2–6.6; Pgs 214-236
3.5 Kinetic Molecular Theory	3.5.A Explain the relationship between the motion of particles and the macroscopic properties of gases with: i. The kinetic molecular theory (KMT). ii. A particulate model. iii. A graphical representation.	3.5.A.1 The kinetic molecular theory (KMT) relates the macroscopic properties of gases to motions of the particles in the gas. The Maxwell-Boltzmann distribution describes the distribution of the kinetic energies of particles at a given temperature.	Ch. 6, Sec. 6.8; Pgs 240-246
		3.5.A.2 All the particles in a sample of matter are in continuous, random motion. The average kinetic energy of a particle is related to its average velocity by the equation: EQN: $KE = \frac{1}{2} mv^2$.	Ch. 6, Sec. 6.8; Pgs 240-246
		3.5.A.3 The Kelvin temperature of a sample of matter is proportional to the average kinetic energy of the particles in the sample.	Ch. 6, Sec. 6.8; Pgs 240-246
		3.5.A.4 The Maxwell-Boltzmann distribution provides a graphical	Ch. 6, Sec. 6.8; Pgs 240-246



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		representation of the energies/velocities of particles at a given temperature.	
3.6 Deviation from Ideal Gas Law	3.6.A Explain the relationship among non-ideal behaviors of gases, interparticle forces, and/or volumes.	3.6.A.1 The ideal gas law does not explain the actual behavior of real gases. Deviations from the ideal gas law may result from interparticle attractions among gas molecules, particularly at conditions that are close to those resulting in condensation. Deviations may also arise from particle volumes, particularly at extremely high pressures.	Ch. 10, Sec. 6.10; Pgs 248-252
3.7 Solutions and Mixtures	3.7.A Calculate the number of solute particles, volume, or molarity of solutions.	3.7.A.1 Solutions, also sometimes called homogeneous mixtures, can be solids, liquids, or gases. In a solution, the macroscopic properties do not vary throughout the sample. In a heterogeneous mixture, the macroscopic properties depend on location in the mixture.	Ch 14, Sec. 14.2, 14.5; Pgs 591-595, 604-610
		3.7.A.2 Solution composition can be expressed in a variety of ways; molarity is the most common method used in the laboratory. EQN: $M = n_{\text{solute}} / L_{\text{solution}}$	Ch 14, Sec. 14.2, 14.5; Pgs 591-595, 604-610
3.8 Representations of Solutions	3.8.A Using particulate models for mixtures: i. Represent interactions between components.	3.8.A.1 Particulate representations of solutions communicate the structure and properties of solutions, by illustration of the relative concentrations of the	Ch 14, Sec. 14.2, 14.4-14.5; Pgs 591-595, 599-610



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	ii. Represent concentrations of components.	components in the solution and/or drawings that show interactions among the components.	
3.9 Separation of Solutions and Mixtures	3.9.A Explain the results of a separation experiment based on intermolecular interactions.	3.9.A.1 The components of a liquid solution cannot be separated by filtration. They can, however, be separated using processes that take advantage of differences in the intermolecular interactions of the components. i. Chromatography (paper, thin-layer, and column) separates chemical species by taking advantage of the differential strength of intermolecular interactions between and among the components of the solution (the mobile phase) and with the surface components of the stationary phase. The resulting chromatogram can be used to infer the relative polarities of components in a mixture. ii. Distillation separates chemical species by taking advantage of the differential strength of intermolecular interactions between and among the components and the effects these interactions have on the vapor pressures of the components in the mixture.	Ch. 1, Sec. 1.3, Pgs 5-8 Chromatography is not explicitly covered in this text
3.10 Solubility	3.10.A Explain the relationship between	3.10.A.1 Substances with similar intermolecular interactions tend to	Ch. 14, Sec. 14.4; Pgs 599-603



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	the solubility of ionic and molecular compounds in aqueous and nonaqueous solvents, and the intermolecular interactions between particles.	be miscible or soluble in one another	
3.11 Spectroscopy and the Electromagnetic Spectrum	3.11.A Explain the relationship between a region of the electromagnetic spectrum and the types of molecular or electronic transitions associated with that region.	3.11.A.1 Differences in absorption or emission of photons in different spectral regions are related to the different types of molecular motion or electronic transition: i. Microwave radiation is associated with transitions in molecular rotational levels. ii. Infrared radiation is associated with transitions in molecular vibrational levels. iii. Ultraviolet/visible radiation is associated with transitions in electronic energy levels.	Ch. 8, Sec. 8.2-8.3; Pgs 318-329
3.12 Properties of Photons	3.12.A Explain the properties of an absorbed or emitted photon in relationship to an electronic transition in an atom or molecule.	3.12.A.1 When a photon is absorbed (or emitted) by an atom or molecule, the energy of the species is increased (or decreased) by an amount equal to the energy of the photon.	Ch. 8, Sec. 8.2; Pgs 318-327
		3.12.A.2 The wavelength of the electromagnetic wave is related to its frequency and the speed of light by the equation:	Not covered in the text thoroughly.



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		EQN: $c = \lambda\nu$. The energy of a photon is related to the frequency of the electromagnetic wave through Planck's equation: EQN: $E = h\nu$	
3.13 Beer-Lambert Law	3.13.A Explain the amount of light absorbed by a solution of molecules or ions in relationship to the concentration, path length, and molar absorptivity	3.13.A.1 The Beer-Lambert law relates the absorption of light by a solution to three variables according to the equation: EQN: $A = \epsilon bc$. The molar absorptivity, ϵ , describes how intensely a chemical species absorbs light of a specific wavelength. The path length, b , and concentration, c , are proportional to the number of light-absorbing particles in the light path.	Not covered in the text thoroughly.
		3.13.A.2 In most experiments the path length and wavelength of light are held constant. In such cases, the absorbance is proportional only to the concentration of absorbing molecules or ions. The spectrophotometer is typically set to the wavelength of maximum absorbance (optimum wavelength) for the species being analyzed to ensure the maximum sensitivity of measurement.	Ch. 8, Sec. 8.4; Pgs 331-335

AP Chemistry Units 4 to 9 correlated to the chapters in Chemistry: A Molecular Approach (6th Edition) by Nivaldo J. Tro.

AP® is a trademark registered and/or owned by the College Board, which was not involved in the production of, and does not endorse, this product



Unit 4: Chemical Reactions

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
4.1 Introduction to Reactions	4.1.A Identify evidence of chemical and physical changes in matter.	4.1.A.1 A physical change occurs when a substance undergoes a change in properties but not a change in composition. Changes in the phase of a substance (solid, liquid, gas) or formation/ separation of mixtures of substances are common physical changes.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202
		4.1.A.2 A chemical change occurs when substances are transformed into new substances, typically with different compositions. Production of heat or light, formation of a gas, formation of a precipitate, and/or color change provide possible evidence that a chemical change has occurred.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202
4.2 Net Ionic Equations	4.2.A Represent changes in matter with a balanced chemical or net ionic equation: i. For physical changes. ii. For given information about the identity of the reactants and/or product. iii. For ions in a given chemical reaction.	4.2.A.1 All physical and chemical processes can be represented symbolically by balanced equations.	Ch. 5, Sec. 5.6; Pgs 185-186
		4.2.A.2 Chemical equations represent chemical changes. These changes are the result of a rearrangement of atoms into new combinations; thus, any representation of a chemical change	Ch. 5, Sec. 5.6; Pgs 185-186



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		must contain equal numbers of atoms of every element before and after the change occurred. Equations thus demonstrate that mass and charge are conserved in chemical reactions.	
		4.2.A.3 Balanced molecular, complete ionic, and net ionic equations are differing symbolic forms used to represent a chemical reaction. The form used to represent the reaction depends on the context in which it is to be used.	Ch. 5, Sec. 5.6; Pgs 185-186
4.3 Representations of Reactions	4.3.A Represent a given chemical reaction or physical process with a consistent particulate model.	4.3.A.1 Balanced chemical equations in their various forms can be translated into symbolic particulate representations.	Ch. 5, Sec. 5.5-5.9; Pgs 181-202
4.4 Physical and Chemical Changes	4.4.A Explain the relationship between macroscopic characteristics and bond interactions for: i. Chemical processes. ii. Physical processes.	4.4.A.1 Processes that involve the breaking and/or formation of chemical bonds are typically classified as chemical processes. Processes that involve only changes in intermolecular interactions, such as phase changes, are typically classified as physical processes.	Ch. 1, Sec. 1.4; Pgs 9-11
		4.4.A.2 Sometimes physical processes involve the breaking of chemical bonds. For example, plausible arguments could be made for the dissolution of a salt in water, as either a physical or chemical process, involves breaking of ionic bonds, and the formation of ion-	Ch. 1, Sec. 1.4; Pgs 9-11



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		dipole interactions between ions and solvent.	
4.5 Stoichiometry	4.5.A Explain changes in the amounts of reactants and products based on the balanced reaction equation for a chemical process.	4.5.A.1 Because atoms must be conserved during a chemical process, it is possible to calculate product amounts by using known reactant amounts, or to calculate reactant amounts given known product amounts.	Ch. 4, Sec. 4.2-4.4; Pgs 141-154
		4.5.A.2 Coefficients of balanced chemical equations contain information regarding the proportionality of the amounts of substances involved in the reaction. These values can be used in chemical calculations involving the mole concept.	Ch. 4, Sec. 4.2-4.4; Pgs 141-154
		4.5.A.3 Stoichiometric calculations can be combined with the ideal gas law and calculations involving molarity to quantitatively study gases and solutions.	Ch. 4, Sec. 4.2-4.4; Pgs 141-154
4.6 Introduction to Titration	4.6.A Identify the equivalence point in a titration based on the amounts of the titrant and analyte, assuming the titration reaction goes to completion.	4.6.A.1 Titrations may be used to determine the amount of an analyte in solution. The titrant has a known concentration of a species that reacts specifically and quantitatively with the analyte. The equivalence point of the titration occurs when the analyte is totally consumed by the reacting species in the titrant. The equivalence point is often indicated by a change in a property (such as color) that occurs when the	Ch. 5, Sec. 5.7, Pgs 187-192



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		equivalence point is reached. This observable event is called the endpoint of the titration.	
4.7 Types of Chemical Reactions	4.7.A Identify a reaction as acid base, oxidation-reduction, or precipitation.	4.7.A.1 Acid-base reactions involve transfer of one or more protons (H^+ ions) between chemical species.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202
		4.7.A.2 Oxidation-reduction (redox) reactions involve transfer of one or more electrons between chemical species, as indicated by changes in oxidation numbers of the involved species. Combustion is an important subclass of oxidation-reduction reactions, in which a species reacts with oxygen gas. In the case of hydrocarbons, carbon dioxide and water are products of complete combustion.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202
		4.7.A.3 In a redox reaction, electrons are transferred from the species that is oxidized to the species that is reduced.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202
		4.7.A.4 Oxidation numbers may be assigned to each of the atoms in the reactants and products; this is often an effective way to identify the oxidized and reduced species in a redox reaction.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		4.7.A.5 Precipitation reactions frequently involve mixing ions in aqueous solution to produce an insoluble or sparingly soluble ionic compound. All sodium, potassium, ammonium, and nitrate salts are soluble in water.	Ch. 4, Sec. 4.5; Pgs 155-157, Ch. 5, Sec. 5.5, 5.7-5.9; Pgs 187-202
4.8 Introduction to Acid-Base 1.B Reactions	4.8.A Identify species as Brønsted Lowry acids, bases, and/or conjugate acid-base pairs, based on proton-transfer involving those species.	4.8.A.1 By definition, a Brønsted-Lowry acid is a proton donor and a Brønsted-Lowry base is a proton acceptor.	Ch. 17, Sec. 17.2-17.3; Pgs 748-752
		4.8.A.2 Only in aqueous solutions, water plays an important role in many acid-base reactions, as its molecular structure allows it to accept protons from and donate protons to dissolved species.	Ch. 17, Sec. 17.2-17.3; Pgs 748-752
		4.8.A.3 When an acid or base ionizes in water, the conjugate acid-base pairs can be identified and their relative strengths compared.	Ch. 17, Sec. 17.2-17.3; Pgs 748-752
4.9 Oxidation-Reduction (Redox) Reactions	4.9.A Represent a balanced redox reaction equation using half-reactions.	4.9.A.1 Balanced chemical equations for redox reactions can be constructed from half-reactions.	Ch. 5, Sec. 5.9; 195-202



Unit 5: Kinetics

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
5.1 Reaction Rates	5.1.A Explain the relationship between the rate of a chemical reaction and experimental parameters	5.1.A.1 The kinetics of a chemical reaction is defined as the rate at which an amount of reactants is converted to products per unit of time.	Ch. 15, Sec. 15.2; Pgs 644-648
		5.1.A.2 The rates of change of reactant and product concentrations are determined by the stoichiometry in the balanced chemical equation.	Ch. 15, Sec. 15.2; Pgs 644-648
		5.1.A.3 The rate of a reaction is influenced by reactant concentrations, temperature, surface area, catalysts, and other environmental factors.	Ch. 15, Sec. 15.2; Pgs 644-648
5.2 Introduction to Rate Law	5.2.A Represent experimental data with a consistent rate law expression.	5.2.A.1 Experimental methods can be used to monitor the amounts of reactants and/or products of a reaction over time and to determine the rate of the reaction.	Ch. 15, Sec. 15.3; Pgs 649-653
		5.2.A.2 The rate law expresses the rate of a reaction as proportional to the concentration of each reactant raised to a power.	Ch. 15, Sec. 15.3,15.4,15.6; Pgs 649-653,668-670
		5.2.A.3 The power of each reactant in the rate law is the order of the reaction with respect to that reactant. The sum of the powers of the reactant concentrations in the rate law is the overall order of the reaction.	Ch. 15, Sec. 15.3; Pgs 649-653



		5.2.A.4 The proportionality constant in the rate law is called the rate constant. The value of this constant is temperature dependent and the units reflect the overall reaction order.	Ch. 15, Sec. 15.3; Pgs 649-653
		5.2.A.5 Comparing initial rates of a reaction is a method to determine the order with respect to each reactant.	Ch. 15, Sec. 15.3; Pgs 649-653
5.3 Concentration Changes Over Time	5.3.A Identify the rate law expression of a chemical reaction using data that show how the concentrations of reaction species change over time.	5.3.A.1 The order of a reaction can be inferred from a graph of concentration of reactant versus time.	Ch. 15, Sec. 15.4; Pgs 654-661
		5.3.A.2 If a reaction is first order with respect to a reactant being monitored, a plot of the natural log (ln) of the reactant concentration as a function of time will be linear.	Ch. 15, Sec. 15.4; Pgs 654-661
		5.3.A.3 If a reaction is second order with respect to a reactant being monitored, a plot of the reciprocal of the concentration of that reactant versus time will be linear.	Ch. 15, Sec. 15.3-15.4; Pgs 652-661
		5.3.A.4 The slopes of the concentration versus time data for zeroth, first, and second order reactions can be used to determine the rate constant for the reaction. Zeroth order: EQN: $[A]_t - [A]_0 = -kt$ First order: EQN: $\ln[A]_t - \ln[A]_0 = -kt$ Second order:	Ch. 15, Sec. 15.4; Pgs 654-661



		EQN: $1/[A]_t - 1/[A]_0 = kt$	
		5.3.A.5 Half-life is a critical parameter for first order reactions because the half-life is constant and related to the rate constant for the reaction by the equation: EQN: $t_{1/2} = 0.693/k$.	Ch. 15, Sec. 15.4; Pgs 654-661
		5.3.A.6 Radioactive decay processes provide an important illustration of first order kinetics.	Ch 21, Sec 21.4-21.6, Pg: 974-983
5.4 Elementary Reactions	5.4.A Represent an elementary reaction as a rate law expression using stoichiometry	5.4.A.1 The rate law of an elementary reaction can be inferred from the stoichiometry of the particles participating in a collision.	Ch. 15, Sec. 15.6; Pgs 668-672
		5.4.A.2 Elementary reactions involving the simultaneous collision of three or more particles are rare.	Not covered in the text thoroughly.
5.5 Collision Model	5.5.A Explain the relationship between the rate of an elementary reaction and the frequency, energy, and orientation of particle collisions	5.5.A.1 For an elementary reaction to successfully produce products, reactants must successfully collide to initiate bond-breaking and bond making events.	Ch. 15, Sec. 15.5; Pgs 662-667
		5.5.A.2 In most reactions, only a small fraction of the collisions leads to a reaction. Successful collisions have both sufficient energy to overcome the activation energy requirements and orientations that allow the bonds to rearrange in the required manner.	Ch. 15, Sec. 15.5; Pgs 662-667
		5.5.A.3 The Maxwell-Boltzmann distribution curve describes the	Ch. 15, Sec. 15.5; Pgs 662-



		distribution of particle energies; this distribution can be used to gain a qualitative estimate of the fraction of collisions with sufficient energy to lead to a reaction, and also how that fraction depends on temperature.	667
5.6 Reaction Energy Profile	5.6.A Represent the activation energy and overall energy change in an elementary reaction using a reaction energy profile.	5.6.A.1 Elementary reactions typically involve the breaking of some bonds and the forming of new ones.	Ch. 15, Sec. 15.5; Pgs 662-667
		5.6.A.2 The reaction coordinate is the axis along which the complex set of motions involved in rearranging reactants to form products can be plotted.	Ch. 15, Sec. 15.5; Pgs 662-667.
		5.6.A.3 The energy profile gives the energy along the reaction coordinate, which typically proceeds from reactants, through a transition state, to products. The energy difference between the reactants and the transition state is the activation energy for the forward reaction.	Ch. 15, Sec. 15.5; Pgs 662-667
		5.6.A.4 The rate of an elementary reaction is temperature dependent because the proportion of particle collisions that are energetic enough to reach the transition state varies with temperature. The Arrhenius equation relates the temperature dependence of the rate of an elementary reaction to the activation energy needed by molecular collisions to reach the	Ch. 15, Sec. 15.5; Pgs 662-667



		transition state.	
5.7 Introduction to Reaction Mechanisms	5.7.A Identify the components of a reaction mechanism.	5.7.A.1 A reaction mechanism consists of a series of elementary reactions, or steps, that occur in sequence. The components may include reactants, intermediates, products, and catalysts.	Ch. 15, Sec. 15.6; Pgs 668-672, Ch. 15, Sec. 15.5; Pgs 662-667
		5.7.A.2 The elementary steps when combined should align with the overall balanced equation of a chemical reaction.	Ch. 15, Sec. 15.5; Pgs 662-667, Ch. 15, Sec. 15.5; Pgs 662-667
		5.7.A.3 A reaction intermediate is produced by some elementary steps and consumed by others, such that it is present only while a reaction is occurring.	Ch. 15, Sec. 15.5; Pgs 662-667, Ch. 15, Sec. 15.6; Pgs 668-672
		5.7.A.4 Experimental detection of a reaction intermediate is a common way to build evidence in support of one reaction mechanism over an alternative mechanism	Ch. 15, Sec. 15.5; Pgs 662-667, Ch. 15, Sec. 15.6; Pgs 668-672
5.8 Reaction Mechanism and Rate Law	5.8.A Identify the rate law for a reaction from a mechanism in which the first step is rate limiting.	5.8.A.1 For reaction mechanisms in which each elementary step is irreversible, or in which the first step is rate limiting, the rate law of the reaction is set by the molecularity of the slowest elementary step (i.e., the rate-limiting step).	Ch. 15, Sec. 15.6; Pgs 668-672
5.9 Pre-Equilibrium Approximation	5.9.A Identify the rate law for a reaction from a mechanism in which the first step is not rate limiting.	5.9.A.1 If the first elementary reaction is not rate limiting, approximations (such as pre-equilibrium) must be made to determine a rate law expression.	Ch. 15, Sec. 15.6; Pgs 668-672



5.10 Multistep Reaction Energy Profile	5.10.A Represent the activation energy and overall energy change in a multistep reaction with a reaction energy profile.	5.10.A.1 Knowledge of the energetics of each elementary reaction in a mechanism allows for the construction of an energy profile for a multistep reaction.	Ch. 15, Sec. 15.6; Pgs 668-672
5.11 Catalysis	5.11.A Explain the relationship between the effect of a catalyst on a reaction and changes in the reaction mechanism	5.11.A.1 In order for a catalyst to increase the rate of a reaction, the addition of the catalyst must increase the number of effective collisions and/ or provide a reaction path with a lower activation energy relative to the original reaction coordinate.	Ch. 15, Sec. 15.7; Pgs 673-678
		5.11.A.2 In a reaction mechanism containing a catalyst, the net concentration of the catalyst is constant. However, the catalyst will frequently be consumed in the rate-determining step of the reaction, only to be regenerated in a subsequent step in the mechanism.	Ch. 15, Sec. 15.7; Pgs 673-678
		5.11.A.3 Some catalysts accelerate a reaction by binding to the reactant(s). The reactants are either oriented more favorably or react with lower activation energy. There is often a new reaction intermediate in which the catalyst is bound to the reactant(s). Many enzymes function in this manner.	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417
		5.11.A.4 Some catalysts involve covalent bonding between the catalyst and the reactant(s). An example is acid-base catalysis, in which a reactant or intermediate either gains or loses a proton. This	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs



		introduces a new reaction intermediate and new elementary reactions involving that intermediate.	402-403, 405-417
		5.11.A.5 In surface catalysis, a reactant or intermediate binds to, or forms a covalent bond with, the surface. This introduces elementary reactions involving these new bound reaction intermediate(s).	Ch. 10, Sec. 10.3-10.4; Pgs 404-411 Ch. 10, Sec. 10.2, 10.4-10.6, 10.11; Pgs 402-403, 405-417

Unit 6: Thermochemistry

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
6.1 Endothermic and Exothermic Processes	6.1.A Explain the relationship between experimental observations and energy changes associated with a chemical or physical transformation	6.1.A.1 Temperature changes in a system indicate energy changes.	Ch. 7, Sec. 7.1-7.3, 7.6; Pgs 267-275, 285-288
		6.1.A.2 Energy changes in a system can be described as endothermic and exothermic processes such as the heating or cooling of a substance, phase changes, or chemical transformations.	Ch. 7, Sec. 7.1-7.3, 7.6; Pgs 267-275, 285-288
		6.1.A.3 When a chemical reaction occurs, the energy of the system either decreases (exothermic reaction), increases (endothermic reaction), or remains the same. For exothermic reactions, the energy lost by the reacting species (system) is gained by the surroundings, as heat transfer from	Ch. 7, Sec. 7.1-7.3, 7.6; Pgs 267-275, 285-288



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		or work done by the system. Likewise, for endothermic reactions, the system gains energy from the surroundings by heat transfer to or work done on the system.	
		6.1.A.4 The formation of a solution may be an exothermic or endothermic process, depending on the relative strengths of intermolecular/interparticle interactions before and after the dissolution process.	Ch. 7, Sec 7.6, Pgs 285-287
6.2 Energy Diagrams	6.2.A Represent a chemical or physical transformation with an energy diagram.	6.2.A.1 A physical or chemical process can be described with an energy diagram that shows the endothermic or exothermic nature of that process.	Ch. 7, Sec. 7.4–7.5; Pgs 276–284
6.3 Heat Transfer and Thermal Equilibrium	6.3.A Explain the relationship between the transfer of thermal energy and	6.3.A.1 The particles in a warmer body have a greater average kinetic energy than those in a cooler body.	Ch. 7, Sec. 7.3; Pgs 271-275
		6.3.A.2 Collisions between particles in thermal contact can result in the transfer of energy. This process is called “heat transfer,” “heat exchange,” or “transfer of energy as heat.”	Ch. 7, Sec. 7.3; Pgs 271-275
		6.3.A.3 Eventually, thermal equilibrium is reached as the particles continue to collide. At thermal equilibrium, the average kinetic energy of both bodies is the same, and hence, their temperatures are the same.	Ch. 7, Sec. 7.3; Pgs 271-275
6.4 Heat Capacity and Calorimetry	6.4.A Calculate the heat q absorbed or released by a system undergoing heating/	6.4.A.1 The heating of a cool body by a warmer body is an important form of energy transfer between two systems. The amount of heat transferred	Ch. 7, Sec. 7.5–7.7; Pgs 282–290



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	cooling based on the amount of the substance, the heat capacity, and the change in temperature	between two bodies may be quantified by the heat transfer equation: EQN: $q = mc\Delta T$. Calorimetry experiments are used to measure the transfer of heat.	
		6.4.A.2 The first law of thermodynamics states that energy is conserved in chemical and physical processes.	Ch 7, Sec 7.3, Pg 271-274
		6.4.A.3 The transfer of a given amount of thermal energy will not produce the same temperature change in equal masses of matter with differing specific heat capacities.	Ch 7, Sec 7.3-7.4, Pg 271-280
		6.4.A.4 Heating a system increases the energy of the system, while cooling a system decreases the energy of the system.	Ch 7, Sec 7.3-7.4, Pg 271-280
		6.4.A.5 The specific heat capacity of a substance and the molar heat capacity are both used in energy calculations.	Ch 7, Sec 7.3-7.4, Pg 271-280
		6.4.A.6 Chemical systems change their energy through three main processes: heating/cooling, phase transitions, and chemical reactions.	Ch 7, Sec 7.3-7.4, Pg 271-280
		6.4.A.7 In calorimetry experiments involving dissolution, temperature changes of the mixture within the calorimeter can be used to determine the direction of energy flow. If the temperature of the mixture increases, thermal energy is released by the dissolution process (exothermic). If the temperature of the mixture decreases, thermal energy is absorbed by the dissolution process (endothermic).	Ch 7, Sec 7.3-7.4, Pg 271-280



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
6.5 Energy of Phase Changes	6.5.A Explain changes in the heat q absorbed or released by a system undergoing a phase transition based on the amount of the substance in moles and the molar enthalpy of the phase transition	6.5.A.1 Energy must be transferred to a system to cause a substance to melt (or boil). The energy of the system therefore increases as the system undergoes a solid-to-liquid (or liquid to-gas) phase transition. Likewise, a system releases energy when it freezes (or condenses). The energy of the system decreases as the system undergoes a liquid-to-solid (or gas to-liquid) phase transition. The temperature of a pure substance remains constant during a phase change.	Ch. 12, Sec. 12.6-12.7; Pgs 530-534
		6.5.A.2 The energy absorbed during a phase change is equal to the energy released during a complementary phase change in the opposite direction. For example, the molar enthalpy of condensation of a substance is equal to the negative of its molar enthalpy of vaporization. Similarly, the molar enthalpy of fusion can be used to calculate the energy absorbed when melting a substance and the energy released when freezing a substance.	Ch. 12, Sec. 12.6-12.7; Pgs 530-534
6.6 Introduction to Enthalpy of Reaction	6.6.A Calculate the heat q absorbed or released by a system undergoing a chemical reaction in relationship to the amount of the reacting substance in moles and the molar enthalpy of reaction.	6.6.A.1 The enthalpy change of a reaction gives the amount of heat energy released (for negative values) or absorbed (for positive values) by a chemical reaction at constant pressure.	Ch. 7, Sec. 7.8; Pgs 290-292



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		6.6.A.2 When the products of a reaction are at a different temperature than their surroundings, they exchange energy with the surroundings to reach thermal equilibrium. Thermal energy is transferred to the surroundings as the reactants convert to products in an exothermic reaction. Thermal energy is transferred from the surroundings as the reactants convert to products in an endothermic reaction.	Ch. 7, Sec. 7.8; Pgs 290-292
		6.6.A.3 The chemical potential energy of the products of a reaction is different from that of the reactants because of the breaking and forming of bonds. The energy difference results in a change in the kinetic energy of the particles, which manifests as a temperature change.	Ch. 7, Sec. 7.8; Pgs 290-292
6.7 Bond Enthalpies	6.7.A Calculate the enthalpy change of a reaction based on the average bond energies of bonds broken and formed in the reaction.	6.7.A.1 During a chemical reaction, bonds are broken and/or formed, and these events change the potential energy of the system.	Ch. 10, Sec. 10.10; Pgs 430-432
		6.7.A.2 The average energy required to break all of the bonds in the reactant molecules can be estimated by adding up the average bond energies of all the bonds in the reactant molecules. Likewise, the average energy released in forming the bonds in the product molecules can be estimated. If the energy released is greater than the energy required, the reaction is	Ch. 10, Sec. 10.10; Pgs 430-432



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		exothermic. If the energy required is greater than the energy released, the reaction is endothermic.	
6.8 Enthalpy of Formation	6.8.A Calculate the enthalpy change for a chemical or physical process based on the standard enthalpies of formation.	6.8.A.1 Tables of standard enthalpies of formation can be used to calculate the standard enthalpies of reactions. EQN: $\Delta H^{\circ}_{\text{reaction}} = \sum \Delta H^{\circ}_{f, \text{products}} - \sum \Delta H^{\circ}_{f, \text{reactants}}$	Ch. 7, Sec. 7.9; Pgs 293-297
6.9 Hess's Law	6.9.A Represent a chemical or physical process as a sequence of steps.	6.9.A.1 Many processes can be broken down into a series of steps. Each step in the series has its own energy change.	Ch 7, Sec. 7.8; Pgs 290-292
	6.9.B Explain the relationship between the enthalpy of a chemical or physical process and the sum of the enthalpies of the individual steps.	6.9.B.1 Because total energy is conserved (first law of thermodynamics), and each individual reaction in a sequence transfers thermal energy to or from the surroundings, the net thermal energy transferred in the sequence will be equal to the sum of the thermal energy transfers in each of the steps. These thermal energy transfers are the result of potential energy changes among the species in the reaction sequence; thus, at constant pressure, the enthalpy change of the overall process is equal to the sum of the enthalpy changes of the individual steps.	Ch 7, Sec 7.3, Pg 271-274
		6.9.B.2 The following are essential principles of Hess's law: i. When a reaction is reversed, the enthalpy change stays constant in	Ch 7, Sec 7.3, Pg 271-274



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		magnitude but becomes reversed in mathematical sign. ii. When a reaction is multiplied by a factor c , the enthalpy change is multiplied by the same factor c . iii. When two (or more) reactions are added to obtain an overall reaction, the individual enthalpy changes of each reaction are added to obtain the net enthalpy change of the overall reaction.	

Unit 7: Equilibrium

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
7.1 Introduction to Equilibrium	7.1.A Explain the relationship between the occurrence of a reversible chemical or physical process, and the establishment of equilibrium, to experimental observations.	7.1.A.1 Many observable processes are reversible. Examples include evaporation and condensation of water, absorption and desorption of a gas, or dissolution and precipitation of a salt. Some important reversible chemical processes include the transfer of protons in acid-base reactions and the transfer of electrons in redox reactions.	Ch. 16, Sec. 16.2; Pgs 699-701
		7.1.A.2 When equilibrium is reached, no observable changes occur in the system. Reactants and products are simultaneously present, and the concentrations or partial pressures of all species remain constant.	Ch. 16, Sec. 16.2-16.9; Pgs 696-730
		7.1.A.3 The equilibrium state is dynamic. The forward and reverse processes	Ch. 16, Sec. 16.2-16.9; Pgs



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		continue to occur at equal rates, resulting in no net observable change.	696-730
		7.1.A.4 Graphs of concentration, partial pressure, or rate of reaction versus time for simple chemical reactions can be used to understand the establishment of chemical equilibrium.	Ch. 16, Sec. 16.2-16.9; Pgs 696-730
7.2 Direction of Reversible Reactions	7.2.A Explain the relationship between the direction in which a reversible reaction proceeds and the relative rates of the forward and reverse reactions.	7.2.A.1 If the rate of the forward reaction is greater than the reverse reaction, then there is a net conversion of reactants to products. If the rate of the reverse reaction is greater than that of the forward reaction, then there is a net conversion of products to reactants. An equilibrium state is reached when these rates are equal.	Ch. 16, Sec. 16.2; Pgs 699-701
7.3 Reaction Quotient and Equilibrium Constant	7.3.A Represent the reaction quotient Q_c or Q_p , for a reversible reaction, and the corresponding equilibrium expressions $K_c = Q_c$ or $K_p = Q_p$	7.3.A.1 The reaction quotient Q_c describes the relative concentrations of reaction species at any time. For gas phase reactions, the reaction quotient may instead be written in terms of partial pressures as Q_p. The reaction quotient tends toward the equilibrium constant such that at equilibrium $K_c = Q_c$ and $K_p = Q_p$. As examples, for the reaction $a A + b B \rightleftharpoons c C + d D$ the law of mass action indicates that the equilibrium expression for (K_c, Q_c) is $\text{EQN: } K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}$ and that for (K_p, Q_p) is $\text{EQN: } K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b}$	Ch. 16, Sec. 16.3-16.5, 16.7; Pgs 702-709, 713-714
		7.3.A.2 The reaction quotient does not include substances whose	Ch. 16, Sec. 16.3-16.5, 16.7;



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		concentrations (or partial pressures) are independent of the amount, such as for solids and pure liquids.	Pgs 702-709, 713-714
7.4 Calculating the Equilibrium Constant	7.4.A Calculate K_c or K_p based on experimental observations of concentrations or pressures at equilibrium.	7.4.A.1 Equilibrium constants can be determined from experimental measurements of the concentrations or partial pressures of the reactants and products at equilibrium.	Ch. 16, Sec. 16.6; Pgs 710-712
7.5 Magnitude of the Equilibrium Constant	7.5.A Explain the relationship between very large or very small values of K and the relative concentrations of chemical species at equilibrium	7.5.A.1 Some equilibrium reactions have very large K values and proceed essentially to completion. Others have very small K values and barely proceed at all.	Ch. 16, Sec. 16.3; Pgs 702-705
7.6 Properties of the Equilibrium Constant	7.6.A Represent a multistep process with an overall equilibrium expression, using the constituent K expressions for each individual reaction.	7.6.A.1 When a reaction is reversed, K is inverted.	Ch. 16, Sec. 16.3; Pgs 702-705
		7.6.A.2 When the stoichiometric coefficients of a reaction are multiplied by a factor c , K is raised to the power c .	Ch. 16, Sec. 16.3; Pgs 702-705
		7.6.A.3 When reactions are added together, the K of the resulting overall reaction is the product of the K 's for the reactions that were summed.	Ch. 16, Sec. 16.3; Pgs 702-705



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		7.6.A.4 Since the expressions for K and Q have identical mathematical forms, all valid algebraic manipulations of K also apply to Q .	Ch. 16, Sec. 16.3; Pgs 702-705
7.7 Calculating Equilibrium Concentrations	7.7.A Identify the concentrations or partial pressures of chemical species at equilibrium based on the initial conditions and the equilibrium constant.	7.7.A.1 The concentrations or partial pressures of species at equilibrium can be predicted given the balanced reaction, initial concentrations, and the appropriate K .	Ch. 16, Sec. 16.8; Pgs 715-724
		7.7.A.2 When $Q < K$, the reaction will proceed with a net consumption of reactants and generation of products. When $Q > K$, the reaction will proceed with a net consumption of products and generation of reactants. When $Q = K$, the system is at dynamic equilibrium; both forward and reverse reactions proceed at the same rate, and the proportion of reactants and products remains constant.	Ch. 16, Sec. 16.8; Pgs 715-724
7.8 Representations of Equilibrium	7.8.A Represent a system undergoing a reversible reaction with a particulate mode	7.8.A.1 Particulate representations can be used to describe the relative numbers of reactant and product particles present prior to and at equilibrium, and the value of the equilibrium constant.	Ch. 16, Sec. 16.5-16.6; Pgs 709-712
7.9 Introduction to Le Châtelier's Principle	7.9.A Identify the response of a system at	7.9.A.1 Le Châtelier's principle can be used to predict the response of a system to stresses such as addition or	Ch. 16, Sec. 16.8; Pgs 715-724



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	equilibrium to an external stress, using Le Châtelier's principle	removal of a chemical species, change in temperature, change in volume/pressure of a gas-phase system, or dilution of a reaction system.	
		7.9.A.2 Le Châtelier's principle can be used to predict the effect that a stress will have on experimentally measurable properties such as pH, temperature, and color of a solution.	Ch. 16, Sec 16.7, 16.9; Pgs 713-714, 725-731
7.10 Reaction Quotient and Le Châtelier's Principle	7.10.A Explain the relationships between Q , K , and the direction in which a reversible reaction will proceed to reach equilibrium.	7.10.A.1 A disturbance to a system at equilibrium causes Q to differ from K , thereby taking the system out of equilibrium. The system responds by bringing Q back into agreement with K , thereby establishing a new equilibrium state.	Ch. 16, Sec 16.7, 16.9; Pgs 713-714, 725-731
		7.10.A.2 Some stresses, such as changes in concentration, cause a change in Q only. A change in temperature causes a change in K . In either case, the concentrations or partial pressures of species redistribute to bring Q and K back into equality.	Ch. 16, Sec 16.7, 16.9; Pgs 713-714, 725-731
7.11 Introduction to Solubility Equilibria	7.11.A Calculate the solubility of a salt based on the value of K_{sp} for the salt.	7.11.A.1 The dissolution of a salt is a reversible process whose extent can be described by K_{sp} , the solubility-product constant.	Ch. 18, Sec. 18.5; Pgs 833-838
		7.11.A.2 The solubility of a substance can be calculated from the K_{sp} for the dissolution process. This relationship can also be used to predict the relative solubility of different substances.	Ch. 18, Sec. 18.5; Pgs 833-838
		7.11.A.3 The solubility rules (see 4.7.A.5) can be quantitatively related to K_{sp} , in	Ch. 18, Sec. 18.5; Pgs 833-



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		which K_{sp} values >1 correspond to soluble salts.	838
		7.11.A.4 The molar solubility of one or more species in a saturated solution can be used to calculate the K_{sp} of a substance.	Ch. 18, Sec. 18.5; Pgs 833-838
7.12 Common-Ion Effect	7.12.A Identify the solubility of a salt, and/or the value of K_{sp} for the salt, based on the concentration of a common ion already present in solution.	7.12.A.1 The solubility of a salt is reduced when it is dissolved into a solution that already contains one of the ions present in the salt. The impact of this “common-ion effect” on solubility can be understood qualitatively using Le Châtelier’s principle or calculated from the K_{sp} for the dissolution process	Ch. 18, Sec. 18.5; Pgs 833-838

Unit 8: Acids and Bases

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
8.1 Introduction to Acids and Bases	8.1.A Calculate the values of pH and pOH, based on K_w and the concentration of all species present in a neutral solution of water	8.1.A.1 The concentrations of hydronium ion and hydroxide ion are often reported as pH and pOH, respectively. EQN: $\text{pH} = -\log[\text{H}_3\text{O}^+]$ EQN: $\text{pOH} = -\log[\text{OH}^-]$ The terms “hydrogen ion” and “hydronium ion” and the symbols $\text{H}^{+(aq)}$ and $\text{H}_3\text{O}^{+(aq)}$ are often used interchangeably for the aqueous ion of hydrogen. Hydronium ion and $\text{H}_3\text{O}^{+(aq)}$ are preferred, but $\text{H}^{+(aq)}$ is also accepted on the AP Exam.	Ch. 17, Sec. 17.1-17.3; Pgs 747-752
		8.1.A.2 Water autoionizes with an equilibrium constant K_w . EQN: $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1.0 \times 10^{-14}$ at	Not covered in the text thoroughly.



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		25°C	
		8.1.A.3 In pure water, $\text{pH} = \text{pOH}$ is called a neutral solution. At 25°C, $\text{p}K_w = 14.0$ and thus $\text{pH} = \text{pOH} = 7.0$. EQN: $\text{p}K_w = 14 = \text{pH} + \text{pOH}$ at 25°C	Not covered in the text thoroughly.
		8.1.A.4 The value of K_w is temperature dependent, so the pH of pure, neutral water will deviate from 7.0 at temperatures other than 25°C.	Not covered in the text thoroughly.
8.2 pH and pOH of Strong Acids and Bases	Calculate pH and pOH based on concentrations of all species in a solution of a strong acid or a strong base.	Molecules of a strong acid (e.g., HCl, HBr, HI, HClO_4 , H_2SO_4 , and HNO_3) will completely ionize in aqueous solution to produce hydronium ions and the conjugate base of the acid. As such, the concentration of H_3O^+ in a strong acid solution is equal to the initial concentration of the strong acid, and thus the pH of the strong acid solution is easily calculated.	Ch. 17, Sec. 17.4-17.7; Pgs 753-773
		8.2.A.2 When dissolved in solution, strong bases (e.g., group I and II hydroxides) completely dissociate to produce hydroxide ions. As such, the concentration of OH^- in a strong base solution is equal to the initial concentration of a group I hydroxide and double the initial concentration of a group II hydroxide, and thus the pOH (and pH) of the strong base solution is easily calculated.	Ch. 17, Sec. 17.4-17.7; Pgs 753-773
8.3 Weak Acid and Base Equilibria	8.3.A Explain the relationship among pH, pOH, and concentrations of all	8.3.A.1 Weak acids react with water to produce hydronium ions. However, only a small percentage of molecules of a weak acid will ionize in this way. Thus,	Ch. 17, Sec. 17.4-17.7; Pgs 753-773



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
	species in a solution of a monoprotic weak acid or weak base.	the concentration of H_3O^+ is much less than the initial concentration of the molecular acid, and the vast majority of the acid molecules remain un-ionized.	
		8.3.A.2 A solution of a weak acid involves equilibrium between an un-ionized acid and its conjugate base. The equilibrium constant for this reaction is K_a, often reported as $\text{p}K_a$. The pH of a weak acid solution can be determined from the initial acid concentration and the $\text{p}K_a$. EQN: $K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$ EQN: $\text{p}K_a = -\log K_a$	Not covered in the text thoroughly.
		8.3.A.3 Weak bases react with water to produce hydroxide ions in solution. However, ordinarily just a small percentage of the molecules of a weak base in solution will ionize in this way. Thus, the concentration of OH^- in the solution does not equal the initial concentration of the base, and the vast majority of the base molecules remain un-ionized	Not covered in the text thoroughly.
		8.3.A.4 A solution of a weak base involves equilibrium between an un-ionized base and its conjugate acid. The equilibrium constant for this reaction is K_b, often reported as $\text{p}K_b$. The pH of a weak base solution can be determined from the initial base concentration and the $\text{p}K_b$.	Ch. 17, Sec. 17.4-17.9; Pgs 753-779
		8.3.A.5 The percent ionization of a weak acid (or base) can be calculated from its $\text{p}K_a$ ($\text{p}K_b$) and the initial concentration of the acid (base). The percent ionization	Ch. 17, Sec. 17.4-17.9; Pgs 753-779



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		can also be calculated from the initial concentration of the acid (base) and the equilibrium concentration of any of the species in the equilibrium expression.	
		8.3.A.6 For any conjugate acid-base pair, the acid ionization constant and base ionization constant are related by K_w : EQN: $K_w = K_a \times K_b$ EQN: $pK_w = pK_a + pK_b$	Ch. 17, Sec. 17.4-17.9; Pgs 753-779
8.4 Acid-Base Reactions and Buffers	8.4.A Explain the relationship among the concentrations of major species in a mixture of weak and strong acids and bases.	8.4.A.1 When a strong acid and a strong base are mixed, they react quantitatively in a reaction represented by the equation: $H^+(aq) + OH^-(aq) \rightarrow H_2O(l)$. The pH of the resulting solution may be determined from the concentration of excess reagent.	Ch. 18, Sec. 18.2; Pgs 804-814
		8.4.A.2 When a weak acid and a strong base are mixed, they react quantitatively in a reaction represented by the equation: $HA(aq) + OH^-(aq) \rightleftharpoons A^-(aq) + H_2O(l)$. If the weak acid is in excess, then a buffer solution is formed, and the pH can be determined from the Henderson-Hasselbalch (H-H) equation (see 8.9.A.1). If the strong base is in excess, then the pH can be determined from the moles of excess hydroxide ion and the total volume of solution. If they are equimolar, then the (slightly basic) pH can be determined from the equilibrium represented by the equation: $A^-(aq) + H_2O(l) \rightleftharpoons HA(aq) + OH^-(aq)$.	Ch. 18, Sec. 18.2; Pgs 804-814



		8.4.A.3 When a weak base and a strong acid are mixed, they will react quantitatively in a reaction represented by the equation: $B(aq) + H_3O^+(aq) \rightleftharpoons HB^+(aq) + H_2O(l).$ If the weak base is in excess, then a buffer solution is formed, and the pH can be determined from the H-H equation. If the strong acid is in excess, then the pH can be determined from the moles of excess hydronium ion and the total volume of solution. If they are equimolar, then the (slightly acidic) pH can be determined from the equilibrium represented by the equation: $HB^+(aq) + H_2O(l) \rightleftharpoons B(aq) + H_3O^+(aq).$	Ch. 18, Sec. 18.2; Pgs 804-814
		8.4.A.4 When a weak acid and a weak base are mixed, they will react to an equilibrium state whose reaction may be represented by the equation: $HA(aq) + B(aq) \rightleftharpoons A^-(aq) + HB^+(aq).$	Ch. 18, Sec. 18.2; Pgs 804-814
8.5 Acid-Base Titrations	8.5.A Explain results from the titration of a mono- or polyprotic acid or base solution, in relation to the properties of the solution and its components.	8.5.A.1 An acid-base reaction can be carried out under controlled conditions in a titration. A titration curve, plotting pH against the volume of titrant added, is useful for summarizing results from a titration.	Ch. 18, Sec. 18.4; Pgs 819-832
		8.5.A.2 At the equivalence point for titrations of monoprotic acids or bases, the number of moles of titrant added is equal to the number of moles of analyte originally present. This relationship can be used to obtain the concentration of the analyte. This is the case for titrations of strong acids/bases and weak acids/bases.	Ch. 18, Sec. 18.4; Pgs 819-832



		8.5.A.3 For titrations of weak acids/bases, it is useful to consider the point halfway to the equivalence point, that is, the half-equivalence point. At this point, there are equal concentrations of each species in the conjugate acid-base pair, for example, for a weak acid $[HA] = [A^-]$. Because $pH = pK_a$ when the conjugate acid and base have equal concentrations, the pK_a can be determined from the pH at the half equivalence point in a titration.	Ch. 18, Sec. 18.4; Pgs 819-832
		8.5.A.4 At the equivalence point, pH is determined by the major species in solution. Strong acid and strong base titrations result in neutral pH at the equivalence point. However, in titrations of weak acids (weak bases), the conjugate base of the weak acid (conjugate acid of the weak base) is present at the equivalence point and can undergo proton-transfer reactions with the surrounding water, producing basic (acidic) solutions.	Ch. 18, Sec. 18.4; Pgs 819-832
		8.5.A.5 For polyprotic acids, titration curves can be used to determine the number of acidic protons. In doing so, the major species present at any point along the curve can be identified, along with the pK_a associated with each proton in a weak polyprotic acid.	Ch. 18, Sec. 18.4; Pgs 819-832
8.6 Molecular Structure and Acids/Bases	8.6.A Explain the relationship between the strength of an acid or base and the structure of the molecule or ion.	8.6.A.1 The protons on a molecule that will participate in acid-base reactions, and the relative strength of these protons, can be inferred from the molecular structure. i. Strong acids (such as HCl, HBr, HI, $HClO_4$, H_2SO_4 weak conjugat, and HNO_3) have very e bases that are stabilized by electronegativity, inductive	Ch. 17, Sec. 17.10; Pgs 786-788



		effects, resonance, or some combination thereof. ii. Carboxylic acids are one common class of weak acid. iii. Strong bases (such as group I and II hydroxides) have very weak conjugate acids. iv. Common weak bases include nitrogenous bases such as ammonia as well as carboxylate ions. v. Electronegative elements tend to stabilize the conjugate base relative to the conjugate acid, and so increase acid strength.	
8.7 pH and pKa	8.7.A Explain the relationship between the predominant form of a weak acid or base in solution at a given pH and the pK_a of the conjugate acid or the pK_b of the conjugate base.	8.7.A.1 The protonation state of an acid or base (i.e., the relative concentrations of HA and A^-) can be predicted by comparing the pH of a solution to the pK_a of the acid in that solution. When solution pH < acid pK_a , the acid form has a higher concentration than the base form. When solution pH > acid pK_a , the base form has a higher concentration than the acid form.	Ch. 18, Sec. 18.2; Pgs 804-814
		8.7.A.2 Acid-base indicators are substances that exhibit different properties (such as color) in their protonated versus deprotonated state, making that property respond to the pH of a solution.	Ch. 18, Sec. 18.2; Pgs 804-814
		8.7.A.3 To ensure accurate results in a titration experiment, acid-base indicators should be selected that have a pK_a close to the pH at the equivalence point	Ch. 18, Sec. 18.2; Pgs 804-814
8.8 Properties of Buffers	8.8.A Explain the relationship between the ability of a buffer to	8.8.A.1 A buffer solution contains a large concentration of both members in a conjugate acid-base pair. The conjugate acid reacts with added base	Ch. 18, Sec. 18.2-18.4; Pgs 804-832



	stabilize pH and the reactions that occur when an acid or a base is added to a buffered solution.	and the conjugate base reacts with added acid. These reactions are responsible for the ability of a buffer to stabilize pH.	
8.9 Henderson-Hasselbalch Equation	8.9.A Identify the pH of a buffer solution based on the identity and concentrations of the conjugate acid-base pair used to create the buffer	8.9.A.1 The pH of the buffer is related to the pK_a the acid and the c of concentration ratio of the conjugate acid-base pair. This relation is a consequence of the equilibrium expression associated with the dissociation of a weak acid, and is described by the Henderson Hasselbalch equation. Adding small amounts of acid or base to a buffered solution does not significantly change the ratio of [A⁻]/[HA] and thus does not significantly change the solutio pH. The change in pH on addition of acid or base to a buffered solution is therefore much less than it would have been in the absence of the buffer. EQN: $\text{pH} = \text{pK}_a + \log \frac{[\text{A}^-]}{[\text{HA}]}$	Ch. 18, Sec. 18.2; Pgs 804-814
8.10 Buffer Capacity	8.10.A Explain the relationship between the buffer capacity of a solution and the relative concentrations of the conjugate acid and conjugate base components of the solution	8.10.A.1 Increasing the concentration of the buffer components (while keeping the ratio of these concentrations constant) keeps the pH of the buffer the same but increases the capacity of the buffer to neutralize added acid or base.	Ch. 18, Sec. 18.2-18.4; Pgs 804-832
		8.10.A.2 When a buffer has more conjugate acid than base, it has a greater buffer capacity for addition of added base than acid. When a buffer has more conjugate base than acid, it	Ch. 18, Sec. 18.2-18.4; Pgs 804-832



		has a greater buffer capacity for addition of added acid than base	
8.11 pH and Solubility	8.11.A Identify the qualitative effect of changes in pH on the solubility of a salt.	8.11.A.1 The solubility of a salt is pH sensitive when one of the constituent ions is a weak acid, a weak base, or the hydroxide ion. These effects can be understood qualitatively using Le Châtelier's principle.	Ch. 18, Sec. 18.5; Pgs 833-838

Unit 9: Thermodynamics and Electrochemistry

AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
9.1 Introduction to Entropy	9.1.A Identify the sign and relative magnitude of the entropy change associated with chemical or physical processes.	9.1.A.1 Entropy increases when matter becomes more dispersed. For example, the phase change from solid to liquid or from liquid to gas results in a dispersal of matter as the individual particles become freer to move and generally occupy a larger volume. Similarly, for a gas, the entropy increases when there is an increase in volume (at constant temperature), and the gas molecules are able to move within a larger space. For reactions involving gas-phase reactants or products, the entropy generally increases when the total number of moles of gas-phase products is greater than the total number of moles of gas-phase reactants.	Ch. 19, Sec. 19.2; Pgs 866-868
		9.1.A.2 Entropy increases when energy is dispersed. According to kinetic molecular theory (KMT), the distribution of kinetic energy among the particles of a gas broadens as the temperature increases. As a result, the entropy of the system increases with	Ch. 19, Sec. 19.2; Pgs 866-868



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		an increase in temperature.	
9.2 Absolute Entropy and Entropy Change	9.2.A Calculate the standard entropy change for a chemical or physical process based on the absolute entropies (standard molar entropies) of the species.	9.2.A.1 The entropy change for a process can be calculated from the absolute entropies of the species involved before and after the process occurs. $\text{EQN: } \Delta S_{\text{reaction}}^{\circ} = \sum S_{\text{products}}^{\circ} - \sum S_{\text{reactants}}^{\circ}$	Ch. 19, Sec. 19.3-19.5, 19.7; Pgs 868-880, 885-888
9.3 Gibbs Free Energy and Thermodynamic Favorability	9.3.A Explain whether a physical or chemical process is thermodynamically favored based on an evaluation of ΔG° .	9.3.A.1 The Gibbs free energy change for a chemical process in which all the reactants and products are present in a standard state (as pure substances, as solutions of 1.0 M concentration, or as gases at a pressure of 1.0 atm (or 1.0 bar)) is given the symbol ΔG° .	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895
		9.3.A.2 The standard Gibbs free energy change for a chemical or physical process is a measure of thermodynamic favorability. Historically, the term “spontaneous” has been used to describe processes for which $\Delta G^{\circ} < 0$. The phrase “thermodynamically favored” is preferred instead so that common misunderstandings (equating “spontaneous” with “suddenly” or “without cause”) can be avoided. When $\Delta G^{\circ} < 0$ for the process, it is said to be thermodynamically favored.	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895
		9.3.A.3 The standard Gibbs free energy change for a physical or chemical process may also be determined from the standard Gibbs free energy of	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference																				
		formation of the reactants and products. EQN: $\Delta G_{\text{reaction}}^{\circ} = \sum \Delta G_{f, \text{products}}^{\circ} - \sum \Delta G_{f, \text{reactants}}^{\circ}$																					
		9.3.A.4 In some cases, it is necessary to consider both enthalpy and entropy to determine if a process will be thermodynamically favored. The freezing of water and the dissolution of sodium nitrate are examples of such phenomena.	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895																				
		9.3.A.5 Knowing the values of ΔH° and ΔS° for a process at a given temperature allows ΔG° to be calculated directly. EQN: $\Delta G^{\circ} = \Delta H^{\circ} - T \Delta S^{\circ}$	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895																				
		In general, the temperature conditions for a process to be thermodynamically favored ($\Delta G^{\circ} < 0$) can be predicted from the signs of ΔH° and ΔS° as shown in the table below: <table border="1"> <thead> <tr> <th>ΔH°</th><th>ΔS°</th><th>Symbols</th><th>$\Delta G^{\circ} < 0$, favored at:</th></tr> </thead> <tbody> <tr> <td>< 0</td><td>> 0</td><td><></td><td>all T</td></tr> <tr> <td>> 0</td><td>< 0</td><td>><</td><td>no T</td></tr> <tr> <td>> 0</td><td>> 0</td><td>>></td><td>high T</td></tr> <tr> <td>< 0</td><td>< 0</td><td><<</td><td>low T</td></tr> </tbody> </table> In cases where $\Delta H^{\circ} < 0$ and $\Delta S^{\circ} > 0$, no calculation of ΔG° is necessary to determine that the process is thermodynamically favored ($\Delta G^{\circ} < 0$). In cases where $\Delta H^{\circ} > 0$ and $\Delta S^{\circ} < 0$, no calculation of ΔG° is necessary to determine that the process is thermodynamically unfavored ($\Delta G^{\circ} > 0$).	ΔH°	ΔS°	Symbols	$\Delta G^{\circ} < 0$, favored at:	< 0	> 0	<>	all T	> 0	< 0	><	no T	> 0	> 0	>>	high T	< 0	< 0	<<	low T	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895
ΔH°	ΔS°	Symbols	$\Delta G^{\circ} < 0$, favored at:																				
< 0	> 0	<>	all T																				
> 0	< 0	><	no T																				
> 0	> 0	>>	high T																				
< 0	< 0	<<	low T																				
9.4 Thermodynamics and Kinetic	9.4.A Explain, in terms of kinetics, why a thermodynamically	9.4.A.1 Many processes that are thermodynamically favored do not occur to any measurable extent, or	Ch. 9, Sec. 19.2; Pgs 866-868																				



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
Control	9.4.A.1 Processes that are thermodynamically favored, but do not occur at a measurable rate, are under “kinetic control.” High activation energy is a common reason for a process to be under kinetic control. The fact that a process does not proceed at a noticeable rate does not mean that the chemical system is at equilibrium. If a process is known to be thermodynamically favored, and yet does not occur at a measurable rate, it is reasonable to conclude that the process is under kinetic control.	they occur at extremely slow rates.	
		9.4.A.2 Processes that are thermodynamically favored, but do not proceed at a measurable rate, are under “kinetic control.” High activation energy is a common reason for a process to be under kinetic control. The fact that a process does not proceed at a noticeable rate does not mean that the chemical system is at equilibrium. If a process is known to be thermodynamically favored, and yet does not occur at a measurable rate, it is reasonable to conclude that the process is under kinetic control.	Ch 7, Sec 7.3, Pg 271-274 Ch. 9, Sec. 19.2; Pgs 866-868
9.5 Free Energy and Equilibrium	9.5.A Explain whether a process is thermodynamically favored using the relationships between K, ΔG°, and T.	9.5.A.1 The phrase “thermodynamically favored” ($\Delta G^\circ < 0$) means that the products are favored at equilibrium ($K > 1$) under standard conditions.	Ch. 19, Sec. 19.10; Pgs 899-901
		9.5.A.2 The equilibrium constant is related to free energy by the equations EQN: $K = e^{-\Delta G^\circ/RT}$ and EQN: $\Delta G^\circ = -RT \ln K$.	Ch. 16, Sec 16.3-16.8, pg 702-717
		9.5.A.3 Connections between K and ΔG° can be made qualitatively through estimation. When ΔG° is near zero, the equilibrium constant will be close to 1. When ΔG° is much larger or much smaller than RT, the value of K deviates strongly from 1.	Ch. 19, Sec. 19.10; Pgs 899-901



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		9.5.A.4 Processes with $\Delta G^\circ < 0$ favor products (i.e., $K > 1$) and those with $\Delta G^\circ > 0$ favor reactants (i.e., $K < 1$).	Ch. 19, Sec. 19.10; Pgs 899-901
9.6 Free Energy of Dissolution	9.6.A Explain the relationship between the solubility of a salt and changes in the enthalpy and entropy that occur in the dissolution process.	9.6.A.1 The free energy change (ΔG°) for dissolution of a substance reflects a number of factors: the breaking of the intermolecular interactions that hold the solid together, the reorganization of the solvent around the dissolved species, and the interaction of the dissolved species with the solvent. It is possible to estimate the sign and relative magnitude of the enthalpic and entropic contributions to each of these factors. However, making predictions for the total change in free energy of dissolution can be challenging due to the cancellations among the free energies associated with the three factors cited.	Ch. 14, Sec. 14.3; Pgs 596-599 Ch. 19, Sec. 19.7-19.8 Pgs 885-895
9.7 Coupled Reactions	9.7.A Explain the relationship between external sources of energy or coupled reactions and their ability to drive thermodynamically unfavorable processes.	9.7.A.1 An external source of energy can be used to make a thermodynamically unfavorable process occur. Examples include: i. Electrical energy to drive an electrolytic cell or charge a battery. ii. Light to drive the overall conversion of carbon dioxide to glucose in photosynthesis.	The text does not specifically cover this topic
		9.7.A. 2. A desired product can be formed by coupling a thermodynamically unfavorable reaction that produces that product to a favorable reaction (e.g., the	The text does not specifically cover this topic



AP Subunit	Enduring Understanding & Learning Objective	Essential Knowledge	Textbook Chapter & Section Reference
		conversion of <i>ATP</i> to <i>ADP</i> in biological systems). In the coupled system, the individual reactions share one or more common intermediates. The sum of the individual reactions produces an overall reaction that achieves the desired outcome and has $\Delta G^\circ < 0$	
9.8 Galvanic (Voltaic) and Electrolytic Cells	9.8.A Explain the relationship between the physical components of an electrochemical cell and the overall operational principles of the cell.	9.8.A.1 Each component of an electrochemical cell (electrodes, solutions in the half-cells, salt bridge, voltage/current measuring device) plays a specific role in the overall functioning of the cell. The operational characteristics of the cell (galvanic vs. electrolytic, direction of electron flow, reactions occurring in each half-cell, change in electrode mass, evolution of a gas at an electrode, ion flow through the salt bridge) can be described at both the macroscopic and particulate levels.	Ch. 20, Sec. 20.3, 20.8; Pgs 919-922, 944-950
		9.8.A.2 Galvanic, sometimes called voltaic, cells involve a thermodynamically favored reaction, whereas electrolytic cells involve a thermodynamically unfavored reaction. Visual representations of galvanic and electrolytic cells are tools of analysis to identify where half-reactions occur and in what direction current flows.	Ch 20, Sec 20.3, pg 919-923
		9.8.A.3 For all electrochemical cells, oxidation occurs at the anode and reduction occurs at the cathode.	Ch 20, Sec 20.3, pg 919-923



9.9 Cell Potential and Free Energy	9.9.A Explain whether an electrochemical cell is thermodynamically favored, based on its standard cell potential and the constituent half-reactions within the cell.	9.9.A.1 Electrochemistry encompasses the study of redox reactions that occur within electrochemical cells. The reactions are either thermodynamically favored (resulting in a positive voltage) or thermodynamically unfavored (resulting in a negative voltage and requiring an externally applied potential for the reaction to proceed).	Ch. 20, Sec. 20.4-20.5; Pgs 923-934
		9.9.A.2 The standard cell potential of electrochemical cells can be calculated by identifying the oxidation and reduction half-reactions and their respective standard reduction potentials.	Ch 20, Sec 20.3, pg 919-923
		9.9.A.3 ΔG° (standard Gibbs free energy change) is proportional to the negative of the cell potential for the redox reaction from which it is constructed. Thus, a cell with a positive E° involves a thermodynamically favored reaction, and a cell with a negative E° involves a thermodynamically unfavored reaction. EQN: $\Delta G^\circ = -nFE^\circ$	Ch. 19, Sec. 19.6, 19.8; Pgs 881-884, 889-895
9.10 Cell Potential Under Nonstandard Conditions	9.10.A Explain the relationship between deviations from standard cell conditions and changes in the cell potential.	9.10.A.1 In a real system under nonstandard conditions, the cell potential will vary depending on the concentrations of the active species. The cell potential is a driving force toward equilibrium; the farther the reaction is from equilibrium, the greater the magnitude of the cell potential.	Ch. 20, Sec. 20.5-20.6; Pgs 931-939
		9.10.A.2 Equilibrium arguments such as Le Châtelier's principle do not apply to electrochemical systems, because the systems are not in equilibrium.	Ch. 16, Sec.16.9; Pgs 931-93



		9.10.A.3 The standard cell potential E° corresponds to the standard conditions of $Q = 1$. As the system approaches equilibrium, the magnitude (i.e., absolute value) of the cell potential decreases, reaching zero at equilibrium (when $Q = K$). Deviations from standard conditions that take the cell further from equilibrium than $Q = 1$ will increase the magnitude of the cell potential relative to E°. Deviations from standard conditions that take the cell closer to equilibrium than $Q = 1$ will decrease the magnitude of the cell potential relative to E°. In concentration cells, the direction of spontaneous electron flow can be determined by considering the direction needed to reach equilibrium.	Not covered in the text thoroughly.
		9.10.A.4 Algorithmic calculations using the Nernst equation are insufficient to demonstrate an understanding of electrochemical cells under nonstandard conditions. However, students should qualitatively understand the effects of concentration on cell potential and use conceptual reasoning, including the qualitative use of the Nernst equation: EQN: $E = E^\circ - (RT/nF) \ln Q$ to solve problems.	Ch. 20, Sec.20.4, pg 935



9.11 Electrolysis and Faraday's Law	9.11.A Calculate the amount of charge flow based on changes in the amounts of reactants and products in an electrochemical cell.	9.11.A.1 Faraday's laws can be used to determine the stoichiometry of the redox reaction occurring in an electrochemical cell with respect to the following: i. Number of electrons transferred ii. Mass of material deposited on or removed from an electrode (as in electroplating) iii. Current iv. Time elapsed v. Charge of ionic species EQN: $I = q/t$	Ch. 20, Sec. 20.7-20.8; Pgs 940-950
-------------------------------------	--	--	-------------------------------------

